

Commentary:

Recruiting brown adipose tissue in human obesity

Paul Trayhurn

Clore Laboratory, University of Buckingham, Buckingham, UK; Obesity Biology Unit,
University of Liverpool, Liverpool, UK; College of Science, King Saud University, Riyadh, Saudi
Arabia

Correspondence should be addressed to:

Professor Paul Trayhurn
Clore Laboratory
University of Buckingham
Hunter Street
Buckingham MK18 1EG
UK

Email: p.trayhurn@liverpool.ac.uk

Fax: +44 1280 820 135

Word count – 985 text (excluding references)

Number of Figures and Tables – 1 Figure

Obesity remains a major biomedical challenge with the associated diseases, particularly insulin resistance and type 2 diabetes, imposing a substantial and increasing burden on healthcare systems. In the US, one-third of adults are classed as obese ($BMI \geq 30$), while in the UK, which has one of the highest incidence rates in Europe, 25% are obese. Conceptually, the treatment of obesity is simple: energy expenditure must exceed energy intake. In the late 1970's it was proposed, primarily from studies on rats and mice, that reduced expenditure on adaptive heat production (thermogenesis) associated with a specialised fat tissue – brown adipose tissue (BAT) – is central to the development of obesity. Correspondingly, for a period stimulation of heat production in BAT was seen as a potential therapeutic route for reversing obesity and there was a search for agents that would stimulate the activity of the tissue.

In the four centuries following its identification different roles were attributed to BAT, but in the early 1960's the tissue was firmly identified as a thermogenic organ generating heat through non-shivering mechanisms. Heat is produced by dissipation of the proton gradient across the mitochondrial inner membrane, resulting in the uncoupling of oxidative phosphorylation (1). This process, critically dependent on the mitochondrial uncoupling protein UCP-1, is stimulated by noradrenaline from the sympathetic innervation acting through β_3 -adrenoceptors (1). The concept that BAT is not only a thermoregulatory organ, but is also implicated in the regulation of energy balance with reduced thermogenesis (cold- and diet-induced) leading to obesity, emerged through studies on obese mice and from overfeeding experiments on rats given a cafeteria diet (2). There was little progress, however, in the application of this perspective to human obesity and interest in BAT waned. This was despite evidence in the 1980s for the presence of UCP-1 and a capacity for the activation of the tissue in adults (3).

A renaissance of interest in BAT was catalysed in the late 2000's following the putative identification of multiple sites of the tissue in adult humans from investigations utilising fluorodeoxyglucose positron emission tomography (FDG-PET) (4). A combination of FDG-PET and the detection of UCP-1 in the same areas resulted in the definitive identification of BAT in many adults (5-6). Further FDG-PET studies demonstrated that BAT activity is inversely related to BMI, being lower in the obese than the lean and declining in older subjects (7-9). Insulin stimulates glucose uptake into BAT in humans and acute cold exposure activates the tissue, as in rodents (6; 10-12). Prolonged intermittent cold exposure, tantamount to cold-acclimation, has been shown to lead to the recruitment of BAT in young, lean subjects (13-15). In contrast, only a limited increase in BAT metabolic activity was evident in patients with type 2 diabetes, these individuals also being overweight and older than those in other studies (16). It has

therefore been unclear whether BAT is inducible in the obese, particularly of an older age, with a resultant enhancement of non-shivering thermogenesis. This is, of course, central to the proposition that BAT is a rational target for the treatment of obesity.

Whether BAT can be recruited in obese humans has been directly addressed in a paper by Hanssen *et al* in this issue of *Diabetes* (17). The authors used FDG-PET to assess glucose uptake into presumptive BAT and white fat depots in a group of obese subjects before and after short-term ‘cold-acclimation’. Cold-acclimation, amounting to exposure to 14–15°C for up to 6 h per day over 10 days, resulted in increased glucose uptake in BAT in 6 of the 10 subjects studied in which the tissue was evident. BAT was also induced by cold-acclimation in a subject where activity was previously absent. As in other studies, BAT activity was inversely related to age, both before and following cold-acclimation. There was, however, no significant relationship with percent body fat (though an inverse trend was evident). In contrast to BAT, cold-acclimation did not increase glucose uptake into subcutaneous and visceral white fat depots implying that ‘browning’ had not occurred (or that brite/beige adipocytes are of limited physiological significance). With skeletal muscle, uptake was significantly increased in the triceps brachii, though not in the scalene muscle, and increased translocation of GLUT4 to the sarcolemma was apparent.

Interestingly, the recruitment of BAT during cold-acclimation was not associated with any elevation in total energy expenditure, nor in non-shivering thermogenesis. This may be a consequence of the modest increases in BAT observed, perhaps reflecting the limited acclimation stimulus that can be imposed on humans, or indicate that the tissue is only a minor contributor to expenditure in adults. The mean BMI of the subjects studied was 32.9, the highest being 36.8, and whether BAT can be recruited in the severely obese is an open question, particularly in the face of tissue hypoxia. Increased glucose uptake in BAT in obese subjects following prolonged cold exposure should impact on glucose homeostasis (plasma glucose is reduced in the cold) and prevent induction of the obesity-associated insulin resistance. Indeed, mouse studies have indicated that brown fat is a major organ in glucose disposal and important in relation to insulin sensitivity, as well as being implicated in triglyceride clearance (18–19).

A key role for BAT in metabolic regulation, particularly during or following cold exposure, suggests that a lack of the tissue/reduction in its activity, could play a role in the metabolic syndrome as well as in obesity (18–19). BAT has certainly re-emerged as a therapeutic target in obesity, and the strategy of utilising selective β_3 -adrenoceptor agonists has met with some recent success (20). Alternative approaches have also been considered, including potentiation of the induction of brite/beige adipocytes, BAT transplantation and stem cell

therapy. Whether the more technically innovative strategies are at all feasible remains unclear. There is also uncertainty whether the expansion and activation of BAT, or browning of white fat depots, could ensure a sustained and sufficient impact on energy balance, and do so without raising cardiovascular concerns.

Duality of interest. No potential conflicts of interest relevant to this article were reported.

References

1. Cannon B, Nedergaard J. Brown adipose tissue: function and physiological significance. *Physiol Rev* 2004;84:277-359
2. Himms-Hagen J. Brown adipose tissue thermogenesis and obesity. *Prog Lipid Res* 1989;28:67-115
3. Lean M, E, J. Brown adipose tissue in humans. *Proc Nutr Soc* 1989;48:243-256
4. Nedergaard J, Bengtsson T, Cannon B. Unexpected evidence for active brown adipose tissue in adult humans. *Am J Physiol Endocrinol Metab* 2007;293:E444-452
5. Cypess AM, Lehman S, Williams G, Tal I, Rodman D, Goldfine AB, Kuo FC, Palmer EL, Tseng YH, Doria A, Kolodny GM, Kahn CR. Identification and importance of brown adipose tissue in adult humans. *N Engl J Med* 2009;360:1509-1517
6. Virtanen KA, Lidell ME, Orava J, Heglind M, Westergren R, Niemi T, Taittonen M, Laine J, Savisto NJ, Enerback S, Nuutila P. Functional brown adipose tissue in healthy adults. *N Engl J Med* 2009;360:1518-1525
7. Pfannenberg C, Werner MK, Ripkens S, Stef I, Deckert A, Schmadl M, Reimold M, Häring H-U, Claussen CD, Stefan N. Impact of age on the relationships of brown adipose tissue with sex and adiposity in humans. *Diabetes* 2010;59:1789-1793
8. Yoneshiro T, Aita S, Matsushita M, Okamatsu-Ogura Y, Kameya T, Kawai Y, Miyagawa M, Tsujisaki M, Saito M. Age-related decrease in cold-activated brown adipose tissue and accumulation of body fat in healthy humans. *Obesity* 2011;19:1755-1760
9. Yoneshiro T, Aita S, Matsushita M, Kameya T, Nakada K, Kawai Y, Saito M. Brown adipose tissue, whole-body energy expenditure, and thermogenesis in healthy adult men. *Obesity* 2011;19:13-16
10. van Marken Lichtenbelt WD, Vanhommerig JW, Smulders NM, Drossaerts JM, Kemerink GJ, Bouvy ND, Schrauwen P, Teule GJ. Cold-activated brown adipose tissue in healthy men. *N Engl J Med* 2009;360:1500-1508
11. Orava J, Nuutila P, Lidell Martin E, Oikonen V, Noponen T, Viljanen T, Scheinin M, Taittonen M, Niemi T, Enerbäck S, Virtanen Kirsi A. Different metabolic responses of human brown adipose tissue to activation by cold and insulin. *Cell Metab* 2011;14:272-279
12. Ouellet V, Labbe SM, Blondin DP, Phoenix S, Guerin B, Haman F, Turcotte EE, Richard D, Carpentier AC. Brown adipose tissue oxidative metabolism contributes to energy expenditure during acute cold exposure in humans. *J Clin Invest* 2012;122:545-552
13. van der Lans AAJJ, Hoeks J, Brans B, Vijgen GHEJ, Visser M, xEb, lle GW, Vosselman MJ, Hansen J, XF, rgensen JA, Wu J, Mottaghay FM, Schrauwen P, van Marken Lichtenbelt WD.

Cold acclimation recruits human brown fat and increases nonshivering thermogenesis. *J Clin Invest* 2013;123:3395-3403

14. Lee P, Smith S, Linderman J, Courville AB, Brychta RJ, Dieckmann W, Werner CD, Chen KY, Celi FS. Temperature-acclimated brown adipose tissue modulates insulin sensitivity in humans. *Diabetes* 2014;63:3686-3698
15. Blondin DP, Labbe SM, Tingelstad HC, Noll C, Kunach M, Phoenix S, Guerin B, Turcotte EE, Carpentier AC, Richard D, Haman F. Increased brown adipose tissue oxidative capacity in cold-acclimated humans. *J Clin Endocrinol Metab* 2014;99:E438-446
16. Hanssen MJW, Hoeks J, Brans B, van der Lans AAJJ, Schaart G, van den Driessche JJ, Jorgensen JA, Boekschedon MV, Hesselink MKC, Havekes B, Kersten S, Mottaghy FM, van Marken Lichtenbelt WD, Schrauwen P. Short-term cold acclimation improves insulin sensitivity in patients with type 2 diabetes mellitus. *Nat Med* 2015;21:863-865
17. Hanssen MJ, van der Lans AA, Brans B, Hoeks J, Jardon KM, Schaart G, Mottaghy FM, Schrauwen P, van Marken Lichtenbelt WD. Short-term cold acclimation recruits brown adipose tissue in obese humans. *Diabetes* 2016; this issue
18. Bartelt A, Bruns OT, Reimer R, Hohenberg H, Ittrich H, Peldschus K, Kaul MG, Tromsdorf UI, Weller H, Waurisch C, Eychmuller A, Gordts PLSM, Rinniger F, Bruegelmann K, Freund B, Nielsen P, Merkel M, Heeren J. Brown adipose tissue activity controls triglyceride clearance. *Nat Med* 2011;17:200-205
19. Stanford KI, Middelbeek RJW, Townsend KL, An D, Nygaard EB, Hitchcox KM, Markan KR, Nakano K, Hirshman MF, Tseng Y-H, Goodyear LJ. Brown adipose tissue regulates glucose homeostasis and insulin sensitivity. *J Clin Invest* 2013;123:215-223
20. Cypess Aaron M, Weiner Lauren S, Roberts-Toler C, Elia Elisa F, Kessler Skyler H, Kahn Peter A, English J, Chatman K, Trauger Sunia A, Doria A, Kolodny Gerald M. Activation of human brown adipose tissue by a β 3-adrenergic receptor agonist. *Cell Metab* 2015;21:33-38

Legend to Figure

Figure 1 - Schematic of the key effects of cold-acclimation on the recruitment of brown adipose tissue. Cell proliferation is stimulated and there are multiple changes within the brown adipocyte, particularly increases in the number of mitochondria and the concentration of UCP1 in the mitochondria. Lipolysis, lipogenesis and glucose uptake are stimulated as part of extensive changes in substrate utilisation and cellular metabolism. SNS, sympathetic nervous system; β_3 , β_3 -adrenoceptor.

Figure 1

