

Post-operative pain management through music-induced analgesia: Investigating musical constructs

Journal:	<i>Psychology of Music</i>
Manuscript ID:	POM-13-548.R4
Manuscript Type:	Original Empirical Investigations
Keyword:	pain, analgesia, distraction, relaxation, harmony, rhythm, cortisol
Abstract:	<p>Distraction and attention-diversion approaches are widely integrated into pain management. Music-induced analgesia, the ability of music to reduce pain perception, is a clinically-relevant approach for managing pain, anxiety and psychological well-being. Research categorises audio-analgesic interventions as homogenous, however enquiry is required to identify which musical constructs may be therapeutically effective. This study investigated the impact of harmony and rhythm on acute, post-operative pain in a sample of 98 patients scheduled for knee surgery. Four music listening groups were compared against controls using silent relaxation. After surgery using standardised anaesthesia, participants undertook a 15-minute intervention per day of in-patient stay. Measures of pain intensity, pain interference, salivary cortisol concentration and mood were obtained. All participants showed reductions in pain from pre- to post-test, indicating silent relaxation was as effective as music listening. Salivary cortisol concentrations showed that music with high harmonicity/rhythmicity reduced cortisol concentration to a greater extent on Day 1 than music with low harmonicity/rhythmicity. These findings validate the homogenous use of auditory distraction for audio-analgesia, and importantly emphasise the core role of compositional musical constructs in maximising early post-operative recovery. Results support the need for additional psychobiological research examining the efficacy of audio-analgesic attention-diversion interventions used in pain management.</p>

Post-operative pain management through audio-analgesia: Investigating musical constructs

Distraction, used for pain management, has become a standardised component of integrated pain management interventions and research (Cepeda, Carr, Lau, & Alvarez, 2006; Eccleston & Crombez, 1999; Eccleston, 2001). In the context of Gate Control Theory and the subsequent Neuromatrix Theory, pain theory asserts that effective distraction can be integral to the redistribution of attentional resources, resulting in pain modulation (Melzack & Wall, 1965; Melzack, 2001). Limited capacity models of attention (Shiffrin & Schneider, 1977) posit that attention is a finite cognitive resource and that the processing of pain signals can be mediated by attentional redirection away from pain and towards absorptive tasks/stimuli (Eccleston, 1994, 1995). Distribution of attentional resources can become flawed when the required task activity exceeds available attentional capacity (Baddeley, 1986; Shiffrin & Schneider, 1977). Hence pain perception can be minimised if the distractor's attentional demands are of sufficient magnitude.

Due to the affective nature of pain, attention-diversion interventions which are emotionally resonant and multi-sensory are salient and effective in engaging attentional resources (Good et al., 2001; MacDonald et al., 2003; McCaul & Malott, 1984). Previous research has considered the use of humour (Mitchell, MacDonald, & Brodie, 2006; Weisenberg, Tepper, & Schwarzwald, 1995; Zweyer, Velker, & Ruch, 2004), relaxation (Cepeda et al., 2006; Evans, 2002a, 2002b; Good et al., 2001, 2008; Good, Anderson, Ahn, Cong, & Stanton-Hicks, 2005; Good, Anderson, Stanton-Hicks, & Makii, 2002), mathematics (Burns, 2006; Mitchell et al., 2006), art (Mitchell, MacDonald, & Knussen, 2008), virtual reality (Chan, Chung, Wong, Lien, & Yang, 2007; Hoffman et al., 2004; Wiederhold & Wiederhold, 2007),

Running Head: Post-operative pain management through audio-analgesia

1
2
3
4
5 video games (Wright & Raudenbush, 2010) and music (Good, 1996; Hekmat & Hertel, 2003;
6
7 McCaffrey & Good, 2000, 2000; McCaul & Malott, 1984), collectively demonstrating that
8
9 pain can be modulated through attention-diversion. Research suggests that music may be
10
11 privileged in a clinical context through its innate emotional resonance, ease of use, universal
12
13 availability, familiarity, non-invasive nature and ease of administration (Lim & Locsin, 2006;
14
15 MacDonald et al., 2003; McCaul & Malott, 1984). Music-induced analgesia, the ability of
16
17 music to attenuate pain perception (MacDonald et al., 2003) is consequently at the forefront
18
19 of clinical attention-diversion research and has been demonstrated intra-operatively (Nilsson,
20
21 Rawal, Unestahl, Zetterberg, & Unosson, 2001), post-operatively (Good et al., 2005; Nilsson,
22
23 Rawal, & Unosson, 2003) in laboratory-induced (Hekmat & Hertel, 2003; Mitchell et al.,
24
25 2006), acute and chronic pain (McCaffrey & Freeman, 2003; Mitchell, MacDonald, Knussen,
26
27 & Serpell, 2007; Zimmerman, Pozehl, Duncan, & Schmitz, 1989).
28
29
30
31

32
33 To date, the efficacy of music in pain management is equivocal; studies have demonstrated
34
35 clinically significant reductions in pain intensity (Good et al., 1999; Heiser, Chiles, Fudge, &
36
37 Gray, 1997; Hekmat & Hertel, 2003; Koch, Kain, Ayoub, & Rosenbaum, 1998), however
38
39 others found that music reduces anxiety and improves psychological well-being (Cadigan et
40
41 al., 2001; MacDonald et al., 2003). If audio-analgesic attention-diversion interventions are to
42
43 be explored and developed further, research is needed to address basic issues regarding the
44
45 intervention. Firstly, what constructs within the sound-source are therapeutically active?
46
47 Music is, in nature, heterogeneous, therefore there may be audio components which
48
49 precipitate greater (or lesser) pain modulation. Research has shown that preference impacts
50
51 upon analgesia, with self-selected music (preferred) or selection from a pre-prepared array
52
53 (quasi-preferred), demonstrating greater pain and anxiety reduction than experimenter-
54
55 selected music (non-preferred) (Hekmat & Hertel, 2003; Mitchell et al., 2006). However,
56
57
58
59
60

Running Head: Post-operative pain management through audio-analgesia

1
2
3
4
5 there has been no research which differentiates constructs *within* music that initiate music-
6
7 induced analgesia. Secondly, music-induced analgesia has predominantly been assessed
8
9 through self-report and minimal research has considered whether this is modelled
10
11 physiologically. Salivary cortisol, a biological marker of stress and anxiety is a potentially
12
13 important indicator of psychobiological change following music-induced analgesia and
14
15 attention-diversion. Music has been shown to halt cortisol elevation after psychological and
16
17 pre-procedural stress (Khalfa, Dalla Bella, Roy, Peretz, & Lupien, 2003; Miluk-Kolasa,
18
19 Obminski, Stupnicki, & Golec, 1994). Research has not considered the effect of music
20
21 listening on cortisol as a marker of post-operative recovery.
22
23
24

25
26 The purpose of this research was to begin to address the questions raised by the current
27
28 heterogenous results in music-induced analgesia research. This study aimed to clarify the
29
30 nature of music-induced analgesia in post-operative pain management following total knee
31
32 arthroplasty (full knee replacement surgery). Musical stimuli were manipulated in terms of
33
34 compositional constructs and standardised pain self-reporting was supplemented by
35
36 assessment of cortisol concentrations.
37
38

39 40 **Method**

41 42 43 *Participants*

44
45 Patients scheduled for primary total knee arthroplasty (primary knee replacement surgery) in
46
47 the Orthopaedics Department, Royal Infirmary of Edinburgh, Scotland, were approached
48
49 regarding participation in the trial. Knee surgery was chosen due to the uniquely high levels
50
51 of pain experienced by patients in their in-patient post-operative recovery days following and
52
53 as a result of the early physiotherapy rehabilitation needed to ensure successful function of
54
55 the prosthesis (Brander et al., 2003).
56
57
58
59
60

Running Head: Post-operative pain management through audio-analgesia

1
2
3
4
5 Participants willing to take part provided written informed consent. Trial recruitment took
6
7 place over 18 months. 122 eligible subjects were identified and 98 recruited. Randomisation
8
9 was according to a computer-generated pseudo-random numerical stratified schedule.

10
11 Patients who could participate in questionnaire-based pain assessment and provide salivary
12
13 cortisol samples were invited to take part. Exclusion criteria included: contraindication to
14
15 central neural blockades, history of allergy to local anaesthetics, non-osteoarthritic chronic
16
17 pain or major hearing deficits (see Figure 1).
18

19
20
21 Insert Figure 1 here: CONSORT Diagram
22

23 24 *Design*

25
26
27 Patients who attained the entry criteria were randomly assigned to one of five experimental
28
29 groups: silent control or four possible types of music listening, using the experimental
30
31 constructs of Harmonicity and Rhythmicity (see Table 1; following Finlay, 2009; Krumhansl,
32
33 2000; Parncutt, 1989). Harmonicity is a global concept reflecting the consonance of music
34
35 through the harmonic series (high + or low -). Rhythmicity is the maintenance of metrical
36
37 (beat) regularity (high + or low -). All clinical staff and participants were blinded to
38
39 allocation. The correct CD was provided by the chief investigator who was, therefore, not
40
41 blinded. Recruiting and post-operative assessment was carried out by the chief investigator
42
43 and trained research nurses.
44
45
46
47

48 All patients undertook a pre-operative assessment at pre-admissions two weeks before
49
50 attendance for surgery. On each day post-surgery all assessment measures were completed
51
52 (Days 1-3). Patient controlled analgesia usage was monitored pre-intervention in the
53
54 immediate 24 hours post-operatively (Day 0).
55
56

57 58 *Music listening intervention*

Running Head: Post-operative pain management through audio-analgesia

1
2
3
4
5 *Musical examples.* A battery of musical examples was created (see Appendix 1). All
6
7 examples were 12–15 minutes in length. 12–15 minutes was deemed appropriate as short-
8
9 term bursts of music listening have been proven effective in an acute pain setting (Lee,
10
11 Chung, Chan, & Chan, 2005; McCaffrey & Good, 2000). This time period of listening was
12
13 selected through interaction with nurses, doctors and consultants who considered it
14
15 appropriate for demonstrating the efficacy of the treatment, yet minimising intrusion into
16
17 clinical care and the daily activity that is required to successfully and efficiently operate an
18
19 orthopaedic ward.
20
21

22
23 Extracts were selected through a pilot study in which musically trained listeners and
24
25 musically untrained listeners were asked to rate 32 tracks for harmonic and rhythmicity.
26
27 Tracks used for rating were across classical, jazz, popular, folk and ethnic music genres to
28
29 encompass all musical tastes. Extracts with the highest (or lowest) ratings by all participants,
30
31 as appropriate to grouping, were included in the final selection for use in this study (see
32
33 Finlay, 2009). Extracts were recorded at a sample rate of 16bits and 44.1kHz. Where multiple
34
35 tracks were used to create a combined length, Pro Tools LE 6.9.2 was used to merge between
36
37 individual extracts. All musical extracts were commercially available, did not include lyrics
38
39 and represented a wide range of possible genres. All record companies and copyright holders
40
41 were contacted and permission to use the music was granted.
42
43
44
45

46 *Audio equipment.* Bose QuietComfort Acoustic Noise Cancelling® headphones that utilise
47
48 full-spectrum noise reduction were used with a Bose portable CD-walkman. The inclusion of
49
50 headphones for research into music and pain has been recommended by Carroll and Seers
51
52 (1998) and Nilsson et al. (2001) in order to reduce environmental noise.
53
54
55
56
57
58
59
60

Running Head: Post-operative pain management through audio-analgesia

Silent Intervention. Around-ear noise cancelling headphones were used to provide a barrier to external sound and to minimise intrusive hospital sound as far as possible.

Pain assessment

This consisted of four structured questionnaires:

Visual Rating Scale (VRS) and Numerical Rating Scale (NRS). Two 11-point subjective assessments of pain intensity were completed for pain at rest (supine on the hospital bed) and on movement (controlled bending of the knee through its full range of motion, see Rakel et al, 2012), using the end-points '0 = no pain' and '10 = pain as bad as you can imagine'.

Results were averaged across both ratings types (NRS and VRS), allowing for inter-correlation and reliability checks between measures (Anderson, 2005).

Short-Form McGill Pain Questionnaire (SFMPQ; Melzack, 1983). The primary outcome measure was expressed as Total Pain Score (TPS) and secondary outcome measures as sensory and affective pain.

Brief Pain Inventory (BPI Short-form; Cleeland, 1992). Results were expressed as a single Mean Pain Interference score.

Profile of Mood States (Profile of Mood States; McNair, Lorr, & Droppleman, 1971). Results were expressed across mood states as a composite Total Mood Disturbance (TMD) score.

Cortisol assessment

Salivary cortisol samples were provided using the passive drool method (Gallagher, Leitch, Massey, McAllister-Williams, & Young, 2006). Samples were analysed for Salivary Cortisol Concentration at time of testing in nmol/l. All saliva samples were frozen immediately on collection and were stored at -20°C. Analysis of spun samples was performed using

Running Head: Post-operative pain management through audio-analgesia

Salimetrics ER Cortisol EIA Kits (Salimetrics, 2006). Cortisol concentration was calculated using an MRX plate reader (Dynex, US).

Peri-operative management

The peri-operative period was from arrival on the ward for surgery until 24 hours afterwards. Previously prescribed analgesics were continued until the day of surgery. Anaesthesia consisted of a spinal anaesthetic with single shot femoral and sciatic nerve blocks. All anaesthesia followed a standardised anaesthetic regimen and therefore was comparable between patients.

Post-operative analgesia was provided using morphine Patient Controlled Analgesic device (PCA; 1mg bolus; 5 minute lockout) overnight on Day 0, and PCA usage was monitored. Post-operative medication and anti-emetics were given as required in accordance with the standardised regimen.

Procedure

Recruitment and Pre-operative Assessment. Patients who were scheduled for primary total knee arthroplasty were identified from the hospital waiting lists. Participants eligible for the study were approached by letter approximately two weeks before their attendance at their Pre-admissions Clinic (PAC).

On PAC arrival, all participants were invited to participate in the study and, to maintain blinding, were told that the research was an investigation into relaxation interventions used during post-operative recovery. If participants wished to continue, informed consent was taken alongside a preliminary assessment, providing baselines for all outcome measures. Subjects were randomised into groups and given the opportunity to (a) listen to one minute

Running Head: Post-operative pain management through audio-analgesia

musical excerpts from the sample CD appropriate to their grouping and to select a track or (b) test the noise-reducing headphones. All participants were informed that they would receive standard care in addition to the relaxation intervention. All patients were requested to refrain from listening to their own music post-operatively and were advised that they could withdraw from the study at any time.

Post-operative assessment. Participants were visited daily at the same time each day and completed the VRS/NRS, SF-MPQ and salivary cortisol sample pre- and post-test. In the interim period, the study apparatus was then set up and music/silence initiated, using headphones attached to a personal CD player. During the 15 minute intervention, participants were asked either sit or lie down, to refrain from any other activity during the intervention session. Curtains were drawn around the hospital bed, visitors were asked to refrain from arriving during the relaxation session and the experimenter left the room to minimise the Hawthorne Effect . The experimenter did not discuss the relaxation with participants before or after the intervention, except as necessary to set up the apparatus. On Days 1 and 3 patients additionally completed the POMS and BPI.

Ethics. The study was approved by NHS Ethics (06/S1101/5) and on-site permissions were granted by the Royal Infirmary of Edinburgh Research and Development Office (2006/R/AN/06). Research was in accordance with the Helsinki Declaration of 1975 (revised 1983).

Statistics: design and analysis

This study was designed as a exploratory prospective single blind randomised trial. The primary outcome measure was pain self-report using the VRS/NRS. Secondary outcome

Running Head: Post-operative pain management through audio-analgesia

measures were Total Pain Score, Mean Pain Interference, Total Mood Disturbance, Cortisol and the sub-scales of the SF-MPQ representing sensory and affective pain. There were no comparable or similar previous randomised controlled trials investigating the effects of musical constructs on post-operative pain. Initial sample size was therefore targeted at 20 per group for this project following (Cohen, 1992). Statistical analysis was carried out using SPSS v.18.

Data from the primary outcome variables was assessed using the Kolmogorov-Smirnov test and Mauchly's test of sphericity. Results were not significant, therefore data was normally distributed and sphericity assumed. Parametric statistical tests are used throughout, with means (standard deviations) presented. Internal consistency was high for all outcome measures (BPI, $\alpha=0.79$; SF-MPQ, $\alpha=0.82$; POMS, $\alpha=0.78$ [55–58]. A minimum alpha level of 0.05 was used for all analyses.

One-way between-group ANOVAs (Group Allocation, 5 levels; control, 4 music types) were computed for the primary outcome measures on all days of testing. Repeated-measures ANOVAs with Bonferroni post-hoc adjustments were computed to determine the impact of Group on Day of Testing (3 levels; post-operative days 1-3) and Time of Testing (2 levels; pre- and post-test). Chi-squared analyses were used to compare proportions. All completed days of testing were used in analysis.

Results

Recruitment characteristics

Of 122 patients approached, 98 consented to take part (80.33%) and 24 did not consent. 89 participants completed the full study and 9 participants completed one or more days of testing (see Fig. 1). Approximately 20 subjects were randomly allocated to each group. 9 participants

withdrew post-randomisation but consented for their complete days of data to be used in analysis.

Of the 98 participants who gave informed consent, 40 were male and 58 were female. Age and arthritic chronicity are reported in Table 2. Men and women were comparable in age at time of surgery (Males $M=68$ years, $SD=7.96$; Females $M=68.12$, $SD=8.14$). All participants suffered from arthritic pain and radiographic arthritis.

Group distribution

A Pearson's chi-square test and a series of one-way ANOVAs were computed to investigate Group Distribution. There was no significant difference between groups in previous experience of arthroplasty surgery ($\chi^2=2.190$, $df=4$, $p=.71$). Similarly, there was no significant difference between Groups in Years of pain and the Age of patients. Groups were therefore comparable in their demographics and surgical histories. Analysis of baseline pain and mood scores and cortisol concentration showed no difference between groups (see Table 2).

Insert Table 2: Demographics and Baseline Pre-admission Scores

Musical background

67.9% ($n=53$) of all experimental group participants did not consider themselves to play a musical instrument, though 32.1% ($n=25$) had played a musical instrument in their lifetime. The most popular musical instrument to play was the piano (17.9%, $n=14$), followed by voice (5.1%, $n=4$). Three participants (3.8%) played folk instruments: two choosing the bagpipes and one playing the accordion. A further five participants had some instrumental experience; two participants played woodwind instruments (2.6%, $n=2$), two played percussion and drum

Running Head: Post-operative pain management through audio-analgesia

1
2
3
4
5 kit (2.6%, $n=2$) and one played the guitar (1.3%, $n=1$). 32% ($n=8$) of participants who played
6
7 a musical instrument had done so for approximately 5–10 years and 24% ($n=6$) had played
8
9 for over 10 years. However, only three participants still played their musical instrument
10
11 (12%, $n=3$). Of the 25 participants who played a musical instrument, 23 (92%) had
12
13 undertaken a minimum of 6 months of Formal Instrumental Musical Tuition (FIMT) on their
14
15 chosen instrument. The majority of formally tutored participants had taken instrumental
16
17 lessons for between 5 and 10 years (43.5%, $n=10$).
18
19

20
21 Univariate between-subjects ANOVAs were conducted to investigate the impact of musical
22
23 experience through Group Allocation. There was no significant effect of length of time
24
25 (years) spent playing a musical instrument or of amount of Formal Instrumental Musical
26
27 Tuition (years). This indicates that the experimental groups were equally matched in their
28
29 distribution of formally trained musicians or participants who had played musical instruments
30
31 in the past.
32
33

34 35 *PCA usage*

36
37
38 Two one-way ANOVAs were computed on the amount of morphine-administered by patients
39
40 on Days 0 and Day 1 post-operatively. There was no difference between groups in the
41
42 amount of morphine used in the PCA on Days 0 or 1. Groups were therefore comparable in
43
44 the rescue analgesic they used post-surgery and pre-intervention. Mean PCA usage on Day 0
45
46 was 15.67 (SD=14.40, median=14) and on Day 1 was 23.62 (SD=21.03, median=18).
47
48

49
50 Significance levels are shown in Table 2.3.5.

51 52 *Gender differences*

Running Head: Post-operative pain management through audio-analgesia

Repeated measures ANOVAs were used to assess the impact of gender on primary and secondary outcome measures. No gender differences were found, indicating that males and females were comparable in all outcome measures.

Primary outcome measures

Insert Table 3: Significance, percentage change and pre- and post-test VRS/NRS scores at rest and on movement (Mean±Standard Deviation)

VRS/NRS scores: rest and movement. As expected, pain on movement was higher and more painful than pain at rest ($p<.0001$), with pain scores on Movement rated as higher and more painful those at rest. All VRS/NRS pre-test scores were higher than post-test scores ($p<.0001$), with a reduction in pain scores across the days of testing ($p<.0001$). Daily significance levels and descriptive statistics are shown in Table 3. There was no significant main effect of group, indicating that pain was not differentially influenced by the type of musical distractor or by the silent relaxation used during the intervention period.

Secondary Outcome Measures

Short-Form McGill Pain Questionnaire: total pain score. Total pain scores improved over the course of the study, with the highest scores on Day 1 and a gradual improvement through to Day 3 ($p<.0001$; see Figure 2). A series of Bonferroni post-hoc comparisons showed that Days 1 and 2 were significantly different from Day 3 ($p<.0001$ and $p=.0304$ respectively), but not from each other. Pre-test scores were significantly higher than post-test scores ($p<.0001$), but again, as there was no impact of group, the type of distractor did not affect total pain scores.

Insert Figure 2 Here

Running Head: Post-operative pain management through audio-analgesia

1
2
3
4
5 *Short-Form McGill Pain Questionnaire: sensory and affective pain.* Both the Sensory and
6
7 Affective Dimensions of Pain displayed pre- to post-test reductions in pain scores (both
8
9 $p < .0001$). For Sensory pain, the magnitude of this reduction was 31% and for Affective pain
10
11 it was 38%. Pain scores across the course of the study showed the same pattern across Days
12
13 of Testing ($p < .0001$) as in previous outcome measures, with Pain at Day 1 highest, with
14
15 significant differences between early days of testing for Sensory pain (Day 1 to Day 2
16
17 $p = .025$; Day 1 to Day 3 $p < .0001$; Day 2 to Day 3 not significant) and later days for Affective
18
19 Pain (Day 1 to Day 2 not significant; Day 1 to Day 3 $p < .0001$; Day 2 to Day 3 $p = .022$).
20
21 Neither dimension showed a significant main effect of Group and there were no further main
22
23 effects or interactions.
24
25

26
27
28 *Brief Pain Inventory: mean pain interference.* The pattern of results differed in terms of
29
30 Mean Pain Interference. Results showed, as before, a significant difference in Days of
31
32 Testing, with Day 1 higher than Day 3 ($p = .0012$). However, there was a main effect of Group
33
34 ($p = .021$) and two-way interaction between Group and Day of Testing ($p = .026$), suggesting
35
36 that the groups did not respond similarly from day-to-day in their MPI scores (see Figure 3
37
38 and Table 4). Where the -- and control groups showed a progressive reduction across the
39
40 three Days of Testing, the ++, +- and -+ groups showed a rise from the PAC to Day 1 and
41
42 decline on Day 3.
43
44

45
46
47 Insert Figure 3 and Table 4 here
48

49
50 *Profile of Mood States: total mood disturbance.* Day of Testing was significant, with an
51
52 improvement in mood disturbance levels as the study progressed ($p < .0001$). Bonferroni post-
53
54 hoc comparisons revealed that the difference between the days was significant at all points:
55
56 scores at the PAC were significantly different from Day 1 ($p < .0001$), and Day 3 ($p = .037$) and
57
58
59
60

Running Head: Post-operative pain management through audio-analgesia

Day 1 was significantly different from Day 3 ($p=.012$). There was no effect of type of intervention (group) on mood disturbance.

Cortisol. Sixty-two participants (63.27%) agreed to provide cortisol samples in their post-operative recovery phase. The remainder of participants chose not to give saliva samples due to: inability to master technique/dry mouth (10.20%), feelings of nausea (8.16%), dislike of spitting (4.08%) or personal choice (13.27%).

Change scores for pre- to post-test cortisol concentrations were computed by subtracting the post-test scores from the pre-test scores for each Day of Testing. Group differences were displayed on Day 1, showing that the type of intervention used impacted upon levels of post-test salivary cortisol ($p=.021$). A series of Bonferroni post-hoc tests comparisons revealed that the ++ group showed a significantly greater reduction from pre- to post-test scores in comparison with the -- group, whose cortisol concentrations rose after the music listening intervention ($p=.019$). Similarly to the other outcome measures, cortisol concentrations reduced across the Days of Testing ($p=.009$), with the highest concentrations demonstrated on Day 1. The three-way interaction between Day of Testing, Time of Testing and Group was significant ($p=.003$). Post-hoc tests revealed that the ++ Group was significantly different from the -- Group on Day 1, but that the groups responded similarly on post-operative days 2 and 3. Descriptive statistics are displayed in Table 5 and Figure 4.

Insert Figure 4 here.

Discussion

This study aimed to identify the therapeutically active properties found within audio attention-diversion for post-operative pain management. Salivary cortisol concentrations were elevated in the immediate post-operative period following the surgical stressor, and

Running Head: Post-operative pain management through audio-analgesia

again reduced by discharge. Pre- to post-test analysis indicated that there were no significant overall reductions in post-test scores, suggesting that the auditory interventions were not successful in reducing endogenous cortisol concentrations in the course of the 15 minute relaxation period. However, post-hoc comparisons on Day 1 demonstrated that pre- to post-test change was significant and greatest in the ++ group and least in the -- group, indicating that the positive/negative combination of both harmonicity and rhythmicity had the greatest impact on the stress response, with harmonic and regular sound minimising cortisol production to the greatest extent. By contrast, music which was low in harmonicity and without rhythmic regularity served to increase cortisol concentrations. The between-groups variability in cortisol concentrations was most strongly evident on the first day post-operatively, implicating the first 24-hours post-operatively as the critical window for audio-analgesic benefit. It also problematises reliance on 'New Age' or 'sedative' (commonly --) music (Phumdoung & Good, 2003; Voss et al., 2004) for relaxation and suggests that ++ music (comparable to the 'pop music' genre) is of greater positive psychobiological impact. Research has demonstrated that fast-paced music elicits greater skin conductance responding than slow-paced music, suggesting that rhythmicity has a potentially activational effect, potentially regulating physiological responses (Carpentier & Potter, 2007). Though there is conflicting evidence, suggesting that pain ratings increase for faster tempi (Kenntner-Mabiala, Gorges, Alpers, Lehmann, & Pauli, 2007). Future research could integrate additional physiological measures and preferred/non-preferred musical preference groupings for greater clarity.

Beyond initial inter-group differences, cortisol concentrations were resistant to pre-test post-test change. The combination of these two findings seem conflicted: between-group divergences suggest music does differentially impact upon cortisol but the lack of post-test

Running Head: Post-operative pain management through audio-analgesia

1
2
3
4
5 change indicates no response. No change is in confirmation of the post-operative pain
6
7 research by Good (2008), but conflicts with pre-procedural or laboratory-induced stress
8
9 research which showed that music did reduce cortisol concentrations in those contexts
10
11 (Khalifa et al., 2003; Miluk-Kolasa et al., 1994). Additionally, it has been shown that
12
13 cognitive-coping strategies could be used to reduce stress and lower cortisol (Hammerfald et
14
15 al., 2006). Current results may be explained by recognising that Total Knee Arthroplasty
16
17 presents a significantly greater clinical stressor than pre-surgical stress, with a more
18
19 pronounced impact on Hypothalamic-Pituitary-Adrenal (HPA) axis activity. The potential of
20
21 relaxation interventions to reduce low-level stress should not be generalised to all possible
22
23 forms of stress, particularly high-level clinical stressors.
24
25
26

27
28 Following surgery, pain was highest on Days 1 and 2 and diminished by Day 3. Magnitude of
29
30 post-test change in the affective pain dimension was greater than that of the sensory
31
32 dimension, suggesting that the interventions had a consistent and significant affective
33
34 psychological impact. The results of this study highlight the importance of affective pain
35
36 management and by extension, recent research has argued that audio-analgesia is dependent
37
38 on personal preference (MacDonald et al., 2003; Mitchell et al., 2008, 2007). The current
39
40 research did exploit preference as participants chose a preferred track from a group-
41
42 appropriate selection, but the positive effects of this could be maximised by recommending
43
44 the use of music from participants' personal collections. In this study, participants allocated
45
46 to the music groups were given the opportunity to select their chosen musical extract from a
47
48 battery of options – essentially employing a quasi-preferred selection method (Finlay, 2009).
49
50 However, the silent control group were given no choice in their use of a non-musical
51
52 relaxation intervention (Finlay & Rogers, 2014). Mitchell & MacDonald (2006) argued for
53
54 the importance of choice in enhancing self-efficacy and perceived control in audio-analgesia,
55
56
57
58
59
60

Running Head: Post-operative pain management through audio-analgesia

1
2
3
4
5 therefore it is possible that it is the variation between intervention preference levels may have
6
7 influenced results. Further research should look at preferred v. quasi-preferred relaxation in
8
9 order to determine whether self-selecting all interventions would further enhance results.
10

11
12 Pain-limited functional interference was minimised as the study progressed. By discharge,
13
14 pain interference levels were below that of pre-admissions, validating the choice to proceed
15
16 with the elective surgical intervention. There was some variability in between-group changes
17
18 over time, with ++, +- and -+ groups showing peaks in functional difficulties on Day 1,
19
20 whereas the silent control and -- group demonstrated progressive, gradual reduction in pain
21
22 interference. Interference was monitored at pre-test only, therefore the discrepancies between
23
24 groups in their improvement in functional ability likely reflects slight differentiations in the
25
26 way in which patients responded to their study and the degree of proprioceptive inhibition
27
28 that occurred immediately following surgery. It is not in response to group allocation as there
29
30 was no between-group main effect. The interventions/placebo responding may have served to
31
32 optimise pain control, which is a primary factor in the speed at which patients return to
33
34 normal function, earlier mobilisation and proficient ambulation (Lingard, Katz, Wright, &
35
36 Sledge, 2004).
37
38
39
40

41
42 This preliminary research found no significant difference between music listening groups or
43
44 silent control in pre- to post-test pain reduction. All groups showed lower post-test scores
45
46 across every measure and dimension of pain assessment. Manipulation of the musical
47
48 constructs of harmonic/rhythmicity did not differentially affect analgesic responses. Quiet
49
50 (silent) relaxation was as effective as music listening in reducing post-operative pain. This
51
52 suggests that constructs *within* an auditory intervention may not alter pain scores. The
53
54 findings validate the inclusion of auditory distraction for post-operative pain management,
55
56 but broadly do not distinguish between or prioritise any single characteristic of auditory
57
58
59
60

Running Head: Post-operative pain management through audio-analgesia

1
2
3
4
5 distraction above another (e.g. harmonicity, rhythmicity, music or silence). This suggests that
6
7 auditory distraction may be a flexible and homogenous intervention in a clinical context,
8
9 providing an opportunity for patients to self-select their attention-diversion strategy. It also
10
11 demonstrates that there is a need to separate and clearly specify differences between ‘music-
12
13 induced analgesia’ and the generic term ‘audio-analgesia’ that is regularly used to refer to
14
15 music-specific interventions (see Mitchell et al, 2006).
16
17

18
19 Results demonstrate a short-term reduction in pain states as a result of the interventions,
20
21 potentially explicable as placebo analgesia. The placebo effect is a stable and documented
22
23 psychobiological phenomenon, whereby “the placebo response is the reduction in a symptom
24
25 as a result of factors relating to a subject’s/patient’s perception of the therapeutic
26
27 intervention” (Vase et al., 2002, p.451). A placebo effect is generally considered to be
28
29 evident in clinical research if there is no differentiation between the control and experimental
30
31 groups (Vase et al., 2002). The phenomenon of placebo analgesia is a robust and validated
32
33 concept and it may play a role in the results of this study, as would be expected in any clinical
34
35 research. However, measures were taken to minimise the impact of placebo through
36
37 controlling the musical extracts chosen, refraining from discussion of the interventions until
38
39 the study conclusion and by blind randomisation of participants.
40
41
42
43

44 It is possible, to argue for an alternative and secondary explanation beyond the placebo effect.
45
46 Noise-reducing headphones were used for the silent control group in order to maintain
47
48 blinding and as a consequence, the audio interventions were therefore compared with a
49
50 baseline treatment as opposed to a (commonly used, unblinded) ‘no treatment’, standard care
51
52 group. Lee et al (2005) found that participants allocated to a headphone control group
53
54 reported that headphones eliminated anxiety occurring due to background noise in the ICU,
55
56 also reducing pain. It is likely therefore that the headphones facilitated separation from
57
58
59
60

external noise, promoted privacy, heightened internal locus of control and reduced noise-related mental fatigue for all participants, regardless of grouping (Evans, 2002b; MacDonald et al., 2003; McCaffrey & Locsin, 2002). This is evidenced in the magnitude of pain score reductions across all groups. The noise-reducing headphones used in this research were therefore an active intervention for all participants, inclusive of controls, which may potentially have minimised between-group comparisons. Future research could integrate dual control groups – headphones and standard care.

The finding that silence was as effective as music reflects the modest results reported in a meta-analysis (Cepeda et al, 2006). It could potentially be argued that in the hospital context, silence functions as a significant auditory distraction of comparable magnitude to music. The hospital environment may be viewed as one in which ‘noise pollution’ is rife. Noise pollution has been defined as an impurity of unpleasant sounds (Cabrera and Lee, 2000). With the beeping of monitoring machines and the activity of staff and other patients, noise levels are often high in hospital. The International Noise Council based with the Environmental Protection Agency in the United States have set guidelines for the noise levels in acute care areas of a hospital (including orthopaedics). Akin to a volume slightly lower than the sound of light traffic or slightly above normal conversation, daytime levels should be approximately 45 dB in the daytime and 20 dB at night (the volume of a quiet conversation) (Bayo et al., 1995). The reality has been proven quite different to the guidelines. Cabrera and Lee (2000) reported that the average noise level of acute care admissions wards at night was recorded at 67 dB.

Music and silence (though noise reducing headphones) amply mask the noise pollution from hospital equipment and activity and can have concomitant psychological, physiological and sociological benefits for the patient (see Finlay, 2013). Though it could be argued that using

Running Head: Post-operative pain management through audio-analgesia

1
2
3
4
5 auditory sources would simply add to the noise pollution, in fact the opposite is true: sound is
6
7 thought to minimise noise pollution (Cabrera and Lee, 2000). Patients in this study
8
9 specifically requested additional use of the interventions for the purposes of sleep-induction,
10
11 and though it was not provided for such uses in this study, in this context, music and silence
12
13 could function as an auditory block in the immediate period before falling asleep. Other
14
15 research has confirmed that sleep quality can impact upon recovery rate. Good et al. (2002)
16
17 investigated pain and sleep quality in gynaecological surgery patients. Those participants who
18
19 had a poor night of sleep had significantly elevated pain during the day following the
20
21 disrupted night. Poor sleep preceded greater pain, but pain did not predict poor sleep. Sleep
22
23 quality was therefore directly proportional to the ensuing levels of pain experienced by
24
25 patients.
26
27
28
29

30
31 It is possible that the pre- and post-test research design may have impacted on the
32
33 significance of results as participants were required to disengage from the stimulus in order to
34
35 complete the questionnaires. The standard protocol in music-induced analgesia research has
36
37 involved the completion of pain scores immediately after a period of music listening (e.g. all
38
39 research by Good et al and MacDonald et al, 2003), as pain ratings require participants to
40
41 divert attention back to pain in order to evaluate their health status. Certainly, the issue of
42
43 durability of results and the most appropriate method of testing has been minimally
44
45 investigated in audio-analgesia literature. Research has shown that music effectively induces
46
47 affective mood changes: Panksepp and Bernatzky (2002) found that music induced desired
48
49 mood changes through positive or negative music. The induced mood persisted at 10 minutes
50
51 post-intervention, but was not statistically evident at 20 minutes. If the results of this study
52
53 were applied to pain, then they would suggest that the post-test SF-MPQ data was taken at
54
55 the peak of the response to the intervention, but the analgesic effects would thereafter have
56
57
58
59
60

1
2
3
4
5 declined quickly, hence they were not reflected in the morning and evening pain ratings.
6
7 Certainly further research is needed to clarify this area. Patients' qualitative perceptions of
8
9 the durability of music-induced analgesic effects have been addressed by Finlay (2013), who
10
11 found heterogeneity in patient views of the longevity of effects, principally related to
12
13 participants' adherence to a strictly biomedical worldview, with a rejection of the potential
14
15 role of psychological stimuli in impacting pain. Future research could consider integrating
16
17 time-delayed repeat assessments to target the length of time across which changes caused by
18
19 psychological interventions are maintained. Alternatively, it could be that there was some
20
21 short-term benefit for all patients through their feelings of involvement in research through
22
23 the Hawthorne Effect (Adair, 1984). Further research should aim to innovate in their study
24
25 designs to clarify this.
26
27
28
29

30 Audio-analgesia and silent relaxation are potentially effective adjunctive treatments for the
31
32 management of post-operative pain. That they were similarly effective demonstrates the
33
34 necessity of a multidisciplinary 'kit' approach to pain interventions (Pellino et al., 2005).
35
36 Cortisol concentrations responded to the interventions most strongly on the first day post-
37
38 operatively, suggesting that attention-diversion strategies may be particularly effective in the
39
40 first 24 hours after surgery and also that the use of positive and auditory interventions should
41
42 be encouraged for patient self-care. This study indicates that further investigation into
43
44 attention-diversion through sound is warranted, but pain scores should be supplemented by
45
46 biological data. The ease of use, desirability and low cost of auditory interventions for
47
48 analgesic purposes argues for their continued use in pain medicine.
49
50
51
52
53
54
55
56
57
58
59
60

References

- Adair, J. G. (1984). The Hawthorne effect: A reconsideration of the methodological artifact. *Journal of Applied Psychology, 69*(2), 334–345. doi:10.1037/0021-9010.69.2.334
- Anderson, K. O. (2005). Role of cutpoints: why grade pain intensity? *Pain, 113*(1-2), 5–6.
doi:10.1016/j.pain.2004.10.024
- Baddeley, A. D. (1986). *Working Memory*. Cambridge: Cambridge University Press.
- Bayo, M. V., Garcia, A. M., and Garcia, A. (1995). Noise levels in an urban hospital and workers' subjective responses. *Arch Environ Health, 50*:247–252.
- Blood, A. J., & Zatorre, R. J. (2001). Intensely pleasurable responses to music correlate with activity in brain regions implicated in reward and emotion. *Proceedings of the National Academy of Sciences of the United States of America, 98*(20), 11818.
- Brander, V. A., Stulberg, S. D., Adams, A. D., Harden, R. N., Bruehl, S., Stanos, S. P., & Houle, T. (2003). Predicting total knee replacement pain: a prospective, observational study. *Clin Orthop Relat Res, 416*, 27–36.
- Burns, J. W. (2006). The role of attentional strategies in moderating links between acute pain induction and subsequent psychological stress: Evidence for symptom-specific reactivity among patients with chronic pain versus healthy nonpatients. *Emotion, 6*(2), 180–192. doi:10.1037/1528-3542.6.2.180
- Cabrera, I. N. and Lee, M. H. M. (2000). Reducing noise pollution in the hospital setting by establishing a department of sound: A survey of recent research on the effects of noise and music in health care. *Preventative Medicine, 30*:339–345.
- Cadigan, M. E., Haldeman, S. M., McNamara, M. E., Noyes, D. A., Spadafora, M. A., & Carroll, D. L. (2001). The effects of music on cardiac patients on bed rest. *Progress in Cardiovascular Nursing, 16*(1), 5–13.

Running Head: Post-operative pain management through audio-analgesia

- 1
2
3
4
5 Carpentier, F. R. D., & Potter, R. F. (2007). Effects of music on physiological arousal:
6
7 Explorations into tempo and genre. *Media Psychology, 10*(3), 339–363.
8
- 9 Carroll, D., & Seers, K. (1998). Relaxation for the relief of chronic pain: a systematic review.
10
11 *Journal of Advanced Nursing, 27*, 476–487.
12
- 13 Cepeda, M. S., Carr, D. B., Lau, J., & Alvarez, H. (2006). Music for pain relief. *Cochrane*
14
15 *Database of Systematic Reviews (Online), (2)*, CD004843.
16
17 doi:10.1002/14651858.CD004843.pub2
18
- 19 Chan, E. A., Chung, J. W. Y., Wong, T. K. S., Lien, A. S. Y., & Yang, J. Y. (2007).
20
21 Application of a virtual reality prototype for pain relief of pediatric burn in Taiwan.
22
23 *Journal of Clinical Nursing, 16*(4), 786–793. doi:10.1111/j.1365-2702.2006.01719.x
24
- 25 Cleeland, C. S. (1992). Brief Pain Inventory (BPI). In *Handbook of Pain Assessment* (pp.
26
27 367–370). Guildford Press.
28
- 29 Cohen, J. (1992). A power primer. *Psychological Bulletin, 112*(1), 155–159.
30
31 doi:10.1037/0033-2909.112.1.155
32
- 33 Eccleston, C. (1994). Chronic pain and attention: a cognitive approach. *British Journal of*
34
35 *Clinical Psychology, 33*, 535–547.
36
- 37 Eccleston, C. (1995). Chronic pain and distraction: an experimental investigation into the role
38
39 of sustained and shifting attention in the processing of chronic persistent pain.
40
41 *Behavioural Research Therapy, 33*(4), 391–405.
42
43
- 44 Eccleston, C. (2001). Role of psychology in pain management. *British Journal of*
45
46 *Anaesthesia, 87*(1), 144–152.
47
- 48 Eccleston, C., & Crombez, G. (1999). Pain Demands Attention: A Cognitive-Affective
49
50 Model of the Interruptive Function of Pain. *Psychological Bulletin, 125*(3), 356–366.
51
52
53
54
55
56
57
58
59
60

Running Head: Post-operative pain management through audio-analgesia

- 1
2
3
4
5 Evans, D. (2002a). Review: Music as a single session intervention reduces anxiety and
6
7 respiratory rate in patients admitted to hospital. *Journal of Advanced Nursing*, 37, 8–
8
9 18.
10
11
12 Evans, D. (2002b). The effectiveness of music as an intervention for hospital patients: a
13
14 systematic review. *Journal of advanced nursing*, 37(1), 8–18.
15
16 Finlay, K. A. (2009). *Audio-analgesia and Multi-disciplinary Pain Management: A*
17
18 *Psychological Investigation into Acute, Post-operative Pain* (PhD thesis). University
19
20 of Edinburgh.
21
22
23 Finlay, K. A. (2013). Familiarity with music in post-operative clinical care. Chapter 4 in
24
25 Daynes, H. & King, E. (Eds.) *Music and Familiarity*.
26
27
28 Finlay, K. A. (2013). Music-induced analgesia in chronic pain: Efficacy and assessment
29
30 through a primary-task paradigm. *Psychology of Music*, 42(3), 325–346.
31
32 doi:10.1177/0305735612471236
33
34
35 Finlay, K. A., & Rogers, J. (2014). Maximizing self-care through familiarity: The role of
36
37 practice effects in enhancing music listening and progressive muscle relaxation for
38
39 pain management. *Psychology of Music*. doi:10.1177/0305735613513311
40
41
42 Gallagher, P., Leitch, M. M., Massey, A. E., McAllister-Williams, R. H., & Young, A. H.
43
44 (2006). Assessing cortisol and dehydroepiandrosterone (DHEA) in saliva: effects of
45
46 collection method. *Journal of Psychopharmacology*, 20(5), 643–649.
47
48
49 Good, M. (1996). Effects of relaxation and music on postoperative pain: a review. *Journal of*
50
51 *Advanced Nursing*, 24, 905–914.
52
53
54 Good, M., Anderson, G. C., Ahn, S., Cong, X., & Stanton-Hicks, M. (2005). Relaxation and
55
56 Music Reduce Pain Following Intestinal Surgery. *Research in Nursing & Health*,
57
58 28(3), 240–251.
59
60

Running Head: Post-operative pain management through audio-analgesia

- 1
2
3
4
5 Good, M., Anderson, G. C., Stanton-Hicks, M., & Makii, M. (2002). Relaxation and Music
6 Reduce Pain After Gynecologic Surgery. *Pain Management Nursing*, 3(2), 61–70.
7
8
9 Good, M., Anderson, G., Wotman, S., Albert, J., Cong, X., Chiang, L., & Bernhofer, E.
10 (2008). (315) Effects of relaxation/music and patient teaching for pain management
11 on salivary cortisol. *The Journal of Pain*, 9(4), 54–54.
12
13
14
15
16 doi:10.1016/j.jpain.2008.01.238
17
18 Good, M., Stanton-Hicks, M., Grass, J. A., Anderson, G. C., Choi, C., Schoolmeesters, L. J.,
19 & Salman, A. (1999). Relief of postoperative pain with jaw relaxation, music and
20 their combination. *Pain*, 81(1-2), 163–172.
21
22
23
24
25 Good, M., Stanton-Hicks, M., Grass, J. A., Anderson, G. C., Lai, H.-L., Roykulcharoen, V.,
26 & Adler, P. A. (2001). Relaxation and music to reduce postsurgical pain. *Journal of*
27
28
29
30
31
32 Hammerfald, K., Eberle, C., Grau, M., Kinsperger, A., Zimmermann, A., Ehlert, U., & Gaab,
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
- J. (2006). Persistent effects of cognitive-behavioral stress management on cortisol responses to acute stress in healthy subjects—a randomized controlled trial. *Psychoneuroendocrinology*, 31(3), 333–339.
- Heiser, R. M., Chiles, K., Fudge, M., & Gray, S. E. (1997). The use of music during the immediate postoperative recovery period. *AORN Journal*, 65(4), 777–785.
- Hekmat, H. M., & Hertel, J. B. (2003). Pain attenuating effects of preferred versus non-preferred musical interventions. *Psychology of Music*, 21, 163–173.
- Hoffman, H. G., Richards, T. L., Coda, B., Bills, A. R., Blough, D., Richards, A. L., & Sharar, S. R. (2004). Modulation of thermal pain-related brain activity with virtual reality: Evidence from fMRI. *NeuroReport: For Rapid Communication of Neuroscience Research*, 15(8), 1245–1248.

Running Head: Post-operative pain management through audio-analgesia

- 1
2
3
4
5 Kenntner-Mabiala, R., Gorges, S., Alpers, G. W., Lehmann, A. C., & Pauli, P. (2007).
6
7 Musically induced arousal affects pain perception in females but not in males: A
8
9 psychophysiological examination. *Biological Psychology*, 75(1), 19–23.
10
11
12 Khalfa, S., Dalla Bella, S., Roy, M., Peretz, I., & Lupien, S. J. (2003). Effects of Relaxing
13
14 Music on Salivary Cortisol Level after Psychological Stress. *Ann N Y Acad Sci*, 999,
15
16 274–376.
17
18 Koch, M. E., Kain, Z. N., Ayoub, C., & Rosenbaum, M. D. (1998). The sedative and
19
20 analgesic sparing effect of music. *Anesthesiology*, 89, 300–306.
21
22
23 Krumhansl, C. L. (2000). Rhythm and pitch in music cognition. *Psychological Bulletin*, 126,
24
25 159–179.
26
27 Lee, O. K. ., Chung, Y. F. ., Chan, M. F., & Chan, W. M. (2005). Music and its effect on the
28
29 physiological responses and anxiety levels of patients receiving mechanical
30
31 ventilation: a pilot study. *Journal of Clinical Nursing*, 14(5), 609–620.
32
33
34 Lim, P. H., & Loesin, R. (2006). Music as nursing intervention for pain in five Asian
35
36 countries. *International nursing review*, 53(3), 189–196.
37
38
39 Lingard, E. A., Katz, J. N., Wright, E. A., & Sledge, C. B. (2004). Predicting the outcome of
40
41 total knee arthroplasty. *The Journal of Bone and Joint Surgery*, 86(10), 2179.
42
43
44 MacDonald, R. A. R., Mitchell, L. A., Dillon, T., Serpell, M. G., Davies, J. B., & Ashley, E.
45
46 A. (2003). An empirical investigation of the anxiolytic and pain reducing effects of
47
48 music. *Psychology of Music*, 31(2), 187–203.
49
50
51 McCaffrey, R., & Freeman, E. (2003). Effect of music on chronic osteoarthritis pain in older
52
53 people. *Journal of Advanced Nursing*, 44(5), 517–524.
54
55
56 McCaffrey, R., & Good, M. (2000). The Lived Experience of Listening to Music While
57
58 Recovering from Surgery. *Journal of Holistic Nursing*, 18(4), 378–390.
59
60

Running Head: Post-operative pain management through audio-analgesia

- 1
2
3
4
5 McCaffrey, R., & Locsin, R. C. (2002). Music Listening as a Nursing Intervention: A
6
7 Symphony of Practice. *Holistic Nursing Practice, 16*(3), 70–77.
8
9 McCaul, K. D., & Malott, J. M. (1984). Distraction and coping with pain. *Psychological*
10
11 *Bulletin, 95*, 516–533.
12
13 McDaniel, L. K., Anderson, K. O., Bradley, L. A., Young, L. D., Turner, R. A., Agudelo, C.
14
15 A., & Keefe, F. J. (1986). Development of an observation method for assessing pain
16
17 behavior in rheumatoid arthritis patients. *Pain, 24*(2), 165–184.
18
19 McNair, D. M., Lorr, M., & Droppleman, L. F. (1971). *Profile of Mood States*. Educational
20
21 and Industrial Testing Service.
22
23
24
25 Melzack, R. (1983). *Pain Measurement and Assessment*. Raven Publishing.
26
27 Melzack, R. (2001). Pain and the Neuromatrix in the Brain. *Journal of Dental Education,*
28
29 *65*(12), 1378–1382.
30
31 Melzack, R., & Wall, P. D. (1965). Pain Mechanisms: a new theory. *Science, 150*, 971–979.
32
33
34 Miluk-Kolasa, B., Obminski, Z., Stupnicki, R., & Golec, L. (1994). Effects of music
35
36 treatment on salivary cortisol in patients exposed to pre-surgical stress. *Experimental*
37
38 *and Clinical Endocrinology, 102*, 118–120.
39
40 Mitchell, L. A., MacDonald, R. A. ., & Knussen, C. (2008). An investigation of the effects of
41
42 music and art on pain perception. *Psychology of Aesthetics, Creativity, and the Arts,*
43
44 *2*(3), 162–170.
45
46 Mitchell, L. A., MacDonald, R. A. R., & Brodie, E. E. (2006). A comparison of the effects of
47
48 preferred music, arithmetic and humour on cold pressor pain. *European journal of*
49
50 *Pain, 10*(4), 343–351.
51
52
53
54
55
56
57
58
59
60

Running Head: Post-operative pain management through audio-analgesia

- 1
2
3
4
5 Mitchell, L. A., MacDonald, R. A. R., Knussen, C., & Serpell, M. G. (2007). A survey
6
7 investigation of the effects of music listening on chronic pain. *Psychology of Music*,
8
9 35(1), 37–57. doi:10.1177/0305735607068887
10
11
12 Nattiez, J. (1990). *Music and Discourse: Toward a Semiology of Music*. Princeton University
13
14 Press.
15
16 Nilsson, U., Rawal, N., Unestahl, L. E., Zetterberg, C., & Unosson, M. (2001). Improved
17
18 recovery after music and therapeutic suggestions during general anaesthesia: a
19
20 double-blind randomised controlled trial. *Acta Anaesthesiologica Scandinavica*, 45,
21
22 812–817.
23
24
25 Nilsson, U., Rawal, N., & Unosson, M. (2003). A comparison of intra-operative or
26
27 postoperative exposure to music—a controlled trial of the effects on postoperative pain.
28
29 *Anaesthesia*, 58(7), 699–703.
30
31
32 Panksepp, J., & Bernatzky, G. (2002). Emotional sounds and the brain: the neuro-affective
33
34 foundations of musical appreciation. *Behavioural Processes*, 60, 133–155.
35
36
37 Parncutt, R. (1989). *Harmony: A Psychoacoustical Approach*. Springer-Verlag.
38
39
40 Pellino, R. A., Gordon, D. B., Engelke, Z. K., Busse, K. L., Collins, M. A., Silver, C. E., &
41
42 Norcross, N. J. (2005). Use of nonpharmacologic interventions for pain and anxiety
43
44 after total hip and total knee arthroplasty. *Orthopaedic Nursing*, 24(3), 182–190.
45
46
47 Phumdoung, S., & Good, M. (2003). Music reduces sensation and distress of labor pain. *Pain*
48
49 *Management Nursing*, 4(2), 54–61.
50
51
52 Price, D. D., Finniss, D. G., and Benedetti, F. (2008). A comprehensive review of the placebo
53
54 effect: Recent advances and current thought. *The Annual Review of Psychology*,
55
56 59:565–590.
57
58
59
60

Running Head: Post-operative pain management through audio-analgesia

- 1
2
3
4
5 Raket, B. A., Blodgett, N. P., Bridget Zimmerman, M., Logsden-Sackett, N., Clark, C.,
6
7 Noiseux, N., Callahan, J., Herr, K., Geasland, K., Yang, X., & Sluka, K. A. (2012).
8
9 Predictors of postoperative movement and resting pain following total knee
10
11 replacement. *Pain*, 153(11), 2192–2203.
12
13 Roy, M., Peretz, I., & Rainville, P. (2008). Emotional valence contributes to music-induced
14
15 analgesia. *Pain*, 134(1-2), 140–147.
16
17 Salimetrics. (2006). Expanded Range High Sensitivity Salivary Cortisol Enzyme
18
19 Immunoassay Kit.
20
21 Shiffrin, R. M., & Schneider, W. (1977). Controlled and automatic human information
22
23 processing II: perceptual learning, automatic attending and a general theory.
24
25 *Psychology Review*, 84, 127–90.
26
27
28
29 Stevenson, C. (1995). Non-pharmacological aspects of acute pain management.
30
31 *Complementary Therapies in Nursing & Midwifery*, 1, 77–84.
32
33 Vase, L., Riley, J. L., and Price, D. D. (2002). A comparison of placebo effects in clinical
34
35 analgesic trials versus studies of placebo analgesia. *Pain*, 99:443–452.
36
37
38 Voss, J. A., Good, M., Yates, B., Baun, M. M., Thompson, A., & Hertzog, M. (2004).
39
40 Sedative music reduces anxiety and pain during chair rest after open-heart surgery.
41
42 *Pain*, 112(1-2), 197–203.
43
44
45 Weisenberg, M., Tepper, I., & Schwarzwald, J. (1995). Humor as a cognitive technique for
46
47 increasing pain tolerance. *Pain*, 63(2), 207–212. doi:10.1016/0304-3959(95)00046-U
48
49 Wiederhold, M. D., & Wiederhold, B. K. (2007). Virtual reality and interactive simulation for
50
51 pain distraction. *Pain Medicine*, 8(Suppl 3), S182–S188. doi:10.1111/j.1526-
52
53 4637.2007.00381.x
54
55
56
57
58
59
60

Running Head: Post-operative pain management through audio-analgesia

1
2
3
4
5 Wright, T., & Raudenbush, B. (2010). Interaction effects of visual distractions, auditory
6
7 distractions and age on pain threshold and tolerance. *North American Journal of*
8
9 *Psychology*, 12(1), 145–158.

10
11 Zimmerman, L., Pozehl, B., Duncan, K., & Schmitz, R. (1989). Effects of music in patients
12
13 who had chronic cancer pain. *Western Journal of Nursing Research*, 11(3), 298–309.

14
15
16 Zweyer, K., Velker, B., & Ruch, W. (2004). Do cheerfulness, exhilaration, and humor
17
18 production moderate pain tolerance? A FACS study. *Humor: International Journal of*
19
20 *Humor Research*, 17(1/2), 85–119.
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 1: CONSORT Diagram

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 2: Total Pain Scores across Time of Testing on the Short-form McGill Pain Questionnaire
215x279mm (72 x 72 DPI)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 3: Mean Pain Interference according to Group Intervention (error bars show ± 1 SD)




Figure 4: Pre- to Post-test change in Cortisol Concentrations. (Positive scores indicate cortisol concentrations improved and *decreased* from pre- to post-test, negative scores indicate cortisol *increased*).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1: Group Allocation

Group	Harmonicity	Rhythmicity
A	+	+
B	-	-
C	-	+
D	+	-
Control	Silent	Silent

For Peer Review

Baseline/Demographics	M±SD	<i>p</i> -value
Age	68.07±8.03	.92
Years of arthritic pain	10.31±10.06	.11
VRS/NRS at Rest	45.96±28.09	.49
VRS/NRS at Movement	66.60±20.81	.35
Mean Pain Interference	5.79±2.03	.74
Total Pain Score	17.11±9.25	.90
Total Mood Disturbance	5.90±27.43	.09
Cortisol (nmol/l)	11.81±13.14	.93
PCA Day 0 (ml)	15.67±14.40	.95
PCA Day 1 (ml)	23.62±21.03	.37

Table 2: Demographics and Baseline Pre-admission Scores

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Day of Testing	Rest				Movement		
	Pre-test	Post-test	% Change	<i>p</i> -value	Pre-test	Post-test	% Change
Day 1	48.68±29.14	43.49±28.43	10.66		73.99±22.43	66.51±24.65	10.11
Day 2	47.78±27.00	38.52±27.28	19.38		65.30±24.41	58.70±28.03	10.11
Day 3	37.12±23.38	30.03±22.00	19.10		60.17±22.64	52.71±23.53	12.39
Overall	44.53±26.51	37.35±25.90	16.38		66.49±23.16	59.31±25.40	10.87
Day of Testing				.0001			
Pre-test to Post-test				.0001			
Movement-Rest				.0005			

Table 3: Significance, percentage change and pre- and post-test VRS/NRS scores at rest and on movement (Mean±Standard Deviation)

Group	Mean Pain Interference				Total Mood Disturbance			
	PAC	Day 1	Day 3	p-value	PAC	Day 1	Day 3	p-value
++	5.81±2.03	6.01±2.97	5.15±2.00		8.93±35.75	24.53±32.58	13.87±25.56	
--	5.31±1.93	7.02±1.73	5.80±2.19		14.67±22.56	25.07±34.18	22.60±34.25	
- +	5.44±2.16	6.85±1.68	6.10±1.64		9.47±24.52	19.93±21.21	10.40±18.09	
+ -	5.97±1.76	6.05±1.83	4.36±2.70		-8.50±21.11	25.44±25.70	14.00±20.18	
Control	6.43±1.98	5.33±1.64	4.94±2.27		15.00±35.75	24.53±32.58	13.87±25.56	
Overall	5.77±1.96	6.25±2.07	5.30±1.19		7.26±28.30	27.23±31.21	17.13±28.81	
Day of Testing				.001				.0001
Between-Groups				.05				.127
2-way Interaction*				.05				.368

Table 4: Significance and Mean Pain Interference and Total Mood Disturbance Scores at Pre-admission (PAC), Day 1 and Day 3 (Mean±Standard Deviation). *Indicates 2-way interaction between Day of Testing and Group

Group	p-value	Day 1			Day 2			Day 3		
		Pre-test	Post-test	% Change	Pre-test	Post-test	% Change	Pre-test	Post-test	% Change
++		33.78±37.68	11.12±6.98	67.08	9.00±4.97	19.79±12.94	-119.89	6.65±5.60	7.68±5.11	-15.49
--		15.59±7.75	27.32±25.97	-75.24	13.36±9.08	13.72±9.36	-2.69	17.70±24.93	14.76±4.94	16.61
-+		17.68±15.84	15.83±11.11	10.46	9.26±4.09	7.96±3.70	14.04	6.68±2.40	7.69±5.77	-15.12
+-		16.73±21.97	19.78±28.87	-18.23	25.96±41.99	21.02±30.24	19.03	8.49±5.19	7.55±3.92	11.07
Control		24.33±25.27	28.38±30.59	-16.65	28.15±29.54	20.21±10.34	28.21	8.28±5.48	9.99±7.91	-20.65
Overall		20.96±22.81	20.84±23.33	0.57	17.67±25.05	16.61±16.98	6.00	9.77±12.49	9.61±6.05	1.64
Day of Testing	.01									
Between-Groups*	.02									
3-way interaction**	.005									

Table 5: Salivary cortisol concentrations before and after music listening or silent control. * Between-groups difference shown only on Day 1. ** 3-way interaction between Day of Testing, Time of Testing and Group.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review

Excerpt No.	Composer/Artist	Title	Excerpt length (mins)
<i>High Harmonicity, Low Rhythmicity (+ -)</i>			
1	Mahler	Symphony 4, Adagietto Sehr Langsam	11.54
2	Vaughan-Williams	Fantasia on a Theme by Thomas Tallis	14.41
3	Glass	Low Symphony, Movt. 1, Subterraneans	15.07
4	Lassus	Hieremiae Propheteiae de Jérémie, Lamentatio Tertia Tertii Diei Missa "Congratulamini mihî": Gloria a 6	11.42
5	Debussy	Prélude à l'après-midi d'un faune	9.02
<i>Low Harmonicity, High Rhythmicity (- +)</i>			
1	Stravinsky	Rite of Spring, Le Sacrifice	18.51
2	Mike Oldfield	Tubular Bells, Part I	15.00
3	Keith Jarrett	(If the) Misfits (Wear it)	13.15
4	Pat Metheny	Sirabhorn Unity Village	14.39
5	Miles Davis	Miles Runs the Voodoo Down	14.01
<i>High Harmonicity, High Rhythmicity (+ +)</i>			
1	Vivaldi	Le Quattro Stagioni, La Primavera	11.16
2	Dvořák	Slavonic Dances, Op. 46, No.s 1, 2, 4	16.18
3	The Rippingtons Jeff Golub Klugh & James	Tourist is Paradise Drop Top Kari	11.47
4	Kartsonakis & Bonar	Vacation in the Sun Return of the Dove Ivory Passage	12.38
5	Trad. arr. Williamson	The Scotch Cap The Lochaben Harper MacGregor's Search The Auld Jew	13.44
6	Stan Getz	I Can't Get Started	11.27
<i>Low Harmonicity, Low Rhythmicity (- -)</i>			
1	Ali Khan & Purna	Emptiness is Form	16.21
2	Pärt	Festina Lente Cantus in Memory of Benjamin Britten	15.26
3	Shakuhachi	Akita No Sugagaki Gekko Roteki	12.42
4	Tommy Smith	Into Silence, No.s 8, 9, 12, 15, 25	14.52